



Abstract

Within the compact central region of ~3% of galaxies, there is evidence for luminous light emission at 22 GHz originating in Microwaves Amplified by Stimulated Emission of Radiation (masers) from water molecules. More than 60% of these detections reveal intensities that are millions of times greater than that of the very first masers discovered in the star-forming spiral arms of our own Milky Way galaxy, and are therefore called megamasers. A fraction of these megamasers are found in a disk-like configuration, offering thus unprecedented tools for **accurate measurements of: (1) direct distances to their host galaxies, independent of assumptions about the geometry of the universe**, as well as **(2) the masses of the central black hole masses that lurk in the centers of these systems, which are usually millions to billions of times heavier than our own Sun**. Unfortunately, there are only a handful of these megamaser disks that we have been able to investigate in great detail. In an attempt to significantly increase the detection rate of these holy grails of astronomy, we are conducting a study of the physical properties of their host galaxies, with the hope of identifying galaxy traits connected to the megamaser disk phenomenon. In this work, we present our techniques for public data collection of the **total flux emitted across the electromagnetic spectrum (i.e., building spectral energy distributions; SEDs) of the host galaxies of all known megamaser disks, with the goal of quantifying the degree to which various energetic components (e.g., black hole accretion, star formation, dust obscuration and reprocessing) contribute to the total galaxy light**. Through SED comparisons of host galaxies that do not host maser emission, along with SED fits of template models from various main emission mechanisms, our SED plots will be used to best diagnose the relations between the 22 GHz emission and that from nuclear accretion, stellar light, or the reprocessing by surrounding dust. **This method will allow more efficient identification of the types of galaxies that are most likely to host megamaser disks in order to increase their detection rate.**


Physics of Masers

MASER – Microwave Amplification by Stimulated Emission of Radiation



Megamasers – 10⁶ times greater than typical galactic masers

Active Galactic Nuclei (Zierr & Biermann 2018)




Megamasers found in a disk-like configuration allow us to:

1. Measure direct distances

- Independent constraint to the Hubble Constant (age of the universe)
- Constrain the geometry of the universe
- Better understand dark energy

2. Measure mass of supermassive black holes


($M_{\text{SMBH}} \sim 10^7 M_{\odot}$).



Spectral Energy Distributions (SED)

SED plots show the energy (νF_{ν}) emitted as a function of frequency (ν)

➤ Fitting multi-frequency observations, i.e. SEDs, attempt to identify the degree to which various energetic components contribute to the total flux of a given galaxy



Components and their emission







- Accretion disk produces X-ray emission
- Gas heated by hot photons from the accretion disk (power law + emission lines)
- Stellar emission from host galaxies peaks in near-IR + contributions from X-ray binaries in soft X-ray
- Hot dust within the torus emits in the mid-IR range
- Dust heated in star formation regions of host galaxy emits in far-IR
- Synchrotron emission, related to the accretion disk and/or radio jets, emits in radio

Importance of SEDs

This study hopes to identify galaxy traits connected to the megamaser disk phenomenon. In particular, by quantifying the light distribution over a large wavelength range, we should be able to address the following questions:


- Are megamaser disks always associated with black hole accretion?
- What mechanisms of the nucleus influence this maser activity?
- Does maser activity require the torus?
- What is an accurate megamaser disk detection rate?

Results: SEDs for best studied megamaser disks



The Megamaser Cosmology Project


- Largest catalog of galaxies surveyed for water maser emission in 22 GHz
- ~3% of all galaxies host maser emission
- ~20% of galaxies that host megamaser emission found in a disk-like configuration
- Previous searches: no systematic analysis of properties of galaxies with maser emissions and those without



To find more water megmaser disks, we must know how to search for them

SED Data: NASA Extragalactic Database (NED)

- NED is a multiwavelength database of for extragalactic objects
- Combines data from large sky surveys and research publications
- Extracted the **frequency (Hz), flux (Jy), uncertainty for the flux measurement, and the aperture sizes** used for each observation



Future Work

- Collect and include **mid-IR photometry** from Wide-field Infrared Survey Explorer (Wright et al. 2010) to the SED plots in order to investigate the link between maser activity and the reprocessing of the nuclear AGN radiation by the surrounding dust (e.g., Stern et al. 2012)
- Proceed with SED fitting to **quantify the contribution of AGN compared to stellar light and other energetic phenomena in these galaxies**

References: Braatz, J., et al., 2009, ApJ, 695, 287; Braatz, et al., 2018, <https://safe.nrao.edu/wiki/bin/view/Main/MegamaserCosmologyProject>; Dattoli, G., Doria, A., et al., 2017, IOP, 4, 1; Herrnstein, et al., 1999, A&A, 20, 165; Hickox, R. & Alexander, D., 2018, ARA&A, 56, 1; Kuo, C., et al., 2010, ApJ, 727, 20; Kuo, C., Constantin, A., et al., 2018, ApJ, 860, 169; Pesce, 2018, PhD Thesis, UVA; Stern, D., et al., 2012, ApJ, 753, 30; Wright E., et al., 2010, AJ, 140, 1868; Zierr, C. & Biermann, P., 2018, A&A, 69, 1

Acknowledgements. This work has been supported by JMU's Physics and Astronomy Department, the 4-VA Collaborative at James Madison University, and the National Science Foundation award NSF:AST #1814594. This research has made use of the NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.